

39 - RELATIONSHIP BETWEEN BMI, SOCIOECONOMIC STATUS, EXERCISE PARTICIPATION AND HABITUAL PHYSICAL ACTIVITY LEVEL OF FATHERS, MOTHERS AND THEIR CHILDREN

ANDRÉ MICHELIN, VILMA PINHEIRO DA C. BRUM, HINAIANA DOS SANTOS MACHADO E WAGNER DE CAMPOS
 Universidade Federal do Paraná Curitiba Paraná Brasil
 andremichelin1@hotmail.com

INTRODUCTION

Overweight is becoming a problem that involves, indiscriminately, people from all social classes, ages, ethnic groups and gender. This disease generates biological consequences, economical, behavioral and social problems. Obesity generates biological complication such as: type II diabetes, insulin resistance, hypertension, orthopedic and vascular problems, chronic inflammation, sleep apnea, and so on (SAELEN & STEPHEN, 2003; VILLARES, RIBEIRO & SILVA, 2003). Economic problems are related to high costs of internment and treatments, the mean medical assistance to these people is more expensive, 36% superior compared to others patients (BUCHALLA, 2003). Behavior difficulties is another problem that includes depression, anxiety, low self-esteem and poor self-image, inclusive during childhood (VILLARES, *et al.*, 2003). Finally, there are social problems that are linked to discrimination in all society and it is believed that the social and economic status can be a predictor of obesity. In some studies, obesity is inversely related to the social status in adult women (LAITINEN, POWER & JÄRVELIN, 2001).

Thinking about overweight and obesity is necessary since childhood because this phase is critical. In the first years of life the metabolism velocity is accelerated and grows new adipocyte. A obese child can have three times the fat cells than a normal child, increasing the possibility of having overweight in the adulthood (GUYTON & HALL, 1997). BAR-OR (2000) estimates that 6-yr-old children have a risk of 50% to became obese adults and for adolescents the risk is bigger (70%). When parents are obese the risk can be higher.

Besides childhood overweight and obesity, there is the diminished physical activity. Leisure activities with low energy expenditure are the most played between children - electronic toys, video game, computer and television have substituted general physical activities. This situation has huge impact in health. Moreover, the high caloric intake nowadays and the genetics (UKKOLA & BOUCHARD, 2002) can generate changes in body mass and BMI in children and adults (BERKEY *et al.*, 2000; ROWLANDS, *et al.*, 1999).

Although children have a higher physical activity level compared to adolescents and adults (OLIVEIRA, 1996), they are more sedentary and obese than children from the past generation. Researchers demonstrated that children, nowadays, are fatter compared to children from 20 years ago (ROSS & PATE, 1987). The situation is more complicated to female; according to some authors (ELIAKIM, *et al.*, 2001; EPSTEIN, *et al.*, 2001; BOREHAM, *et al.*, 1997) there are significant differences in the amount and intensity of physical activities performed by boys and girls, in which males are more active.

Many authors (RIBEIRO, 2001; BLAAK, WESTERTERP, BAR-OR, WOUTERS & SARIS, 1992), indicate that obesity and morbidity related diseases in children and adolescents are increased by sedentary behavior.

Thus, because physical activity is directly related to the body healthy development and growth, it is necessary that children acquire an active lifestyle and take part of exercise programs to guarantee benefits such as psychological, emotional, social improvement, motor abilities and physical fitness (BAR-OR, 2000; OLIVEIRA, 1996).

The initiative of an active behavior should be taken at the family environment. Parents being active increase the possibility of children being active as well. A research by Moore (1991) showed that children of active mothers are twice more active than the ones of inactive mothers and when both parents are active, a child has 5,8 time more chances to be an active child than their count partners with inactive parents.

The relationship between parents and sons is showed also in issues related to obesity and energy expenditure and there is a variety of results to this relationship. Wurmser, *et al* (1998), cited by Treuth, Butte and Wong, (2000) showed that prepubescent girls that have both parents obese presented low energy expenditure. However, Treuth and colleges (2000) determine the energy expenditure through calorimetric method and double labeled water in 8-9 year old girls divided in three groups: both slim parents' girls, both obese parents' girls and one obese and the other slim parents' girls. The results of this study concluded that there were no significant differences in the energy expenditure among the three groups and the greater tendency to obesity is not related to the energy expenditure in girls.

There seems to be a constant integration between child and their parents in the lifestyle (MOORE, 1991) and in the nutritional status, according to ENGSTROM & ANJOS (1996), but in studies with the same objective are rare and this difficult the elaboration of programs to prevent sedentary behavior and obesity in childhood. Thus, the purpose of this study is to make a diagnostic if there is a relationship among BMI, socioeconomic status, exercise practice and parents' habitual physical activity level in schoolchildren aged 7-8 years from particular school system from Curitiba/PR/Brazil.

METHODS

Sample

Fifteen 7-to 8-yr-old schoolchildren of both genders and their parents (15 mothers and 15 fathers) participated in the study. The initial sample was composed by 90 children, however only 15 meet all the criteria: being health, live with both parents, be classified as active or sedentary throughout the physical activity level, parents sign the consent form, parents agree to participate in the study. Written informed consent was obtained form parents. This study was approved by the University of Paraná Human Research Ethics Committee.

Measurement and procedures

Body mass index (BMI)

At first the body mass and stature were collected. Subjects wore comfortable clothes and no shoes and stand in the orthostatic position. Body mass was measured with "FILIZOLA" balance with an accuracy of 100g, for stature it was used a commercial stadiometer. After that, the BMI (kg/m^2) was calculated.

Physical activity level and exercise practice

To assess the physical activity level (PAL) that means all activities performed during the subjects' routine, the three day diary proposed by Bouchard *et al* (1983) was used and modified as an interview format (MACHADO, 2002). Throughout two days of the week and one day of the weekend this diary was possible to obtain the total energy expenditure. This instrument has validity and reliable coefficient above 85%. After calculated the PAL, children were subdivided in two groups: sedentary and active groups. The first one is composed of subjects in which the PAL is less than $39,80 \text{ Kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ and the second one is composed of subjects in which PAL is above $40,50 \text{ Kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$. This classification was established by statistical analyses with more than 200 schoolchildren in previous studies (BRUM, 2004).

To assess the parents' exercise practice, a survey was used in which parents answer if they regularly exercise (at least twice a week and 120 minutes per week) or not. If the answer was positive they were codified as 1 and if not it was codified as 2.

Socioeconomic status

A survey was applied to establish the socioeconomic status based in the ANEP (IBGE 2001). This instrument classifies the families in seven social classes, the higher to the poorer using letters (A1, A2, B1, B2, C, D, E), according to the scholar status of the father, the mother and the families' goods (for example: television, bathroom, servant, etc.). To standardization in the statistic program the social classes were codified in a descendent way (7=A, 6=A2, 5=B1, 4=B2, 3=C, 2=D, 1=E.1). The monetary income was also included.

Statistical analysis

A one-way ANOVA was performed to assess the differences among the variables related to the children (BMI, socioeconomic status, familiar income, exercise practice and parents' PAL). Pearson's product moment correlation was used to determine the relationship among all parameters. A significance level of $p < 0,05$ was selected.

RESULTS AND DISCUSSION

There were no significant differences in the comparison among familiar income, social status, BMI, exercise practice and physical activity level habitual between sedentary and active children (Tables 1 and 2).

Table 1: Values of age, familiar income, social status, BMI, % of parents that exercise and physical activity level for sedentary children.

	Mean	Minimum value	Maximal value	Standard-deviation
Child age (years)	8,15	7,27	9,41	0,63
Family income (# minimum wages)	17,88	10,00	21,00	4,88
Social status	5,50	4,00	6,00	0,76
Father's BMI (kg.m^{-2})	26,47	21,71	31,92	3,30
Mother's BMI (kg.m^{-2})	21,79	18,99	27,34	2,70
Child's BMI (kg.m^{-2})	15,60	11,59	19,67	2,65
Father's exercise (%)	62,5	-	-	-
Mother's exercise (%)	75,0	-	-	-
Father's PAL ($\text{Kcal.kg}^{-1}.\text{day}^{-1}$)	40,87	34,62	54,91	6,73
Mother's PAL ($\text{Kcal.kg}^{-1}.\text{day}^{-1}$)	44,00	38,47	49,82	3,77
Child's PAL ($\text{Kcal.kg}^{-1}.\text{day}^{-1}$)	37,31	35,29	39,73	1,31

Table 2: Values of age, familiar income, social status, BMI, % of parents that exercise and physical activity level for active children

	Média	Mínimo	Máximo	Desvio padrão
Child age (years)	8,32	7,87	8,68	0,28
Family income (# minimum wages)	14,14	7,00	20,00	5,37
Social status	5,14	3,00	7,00	1,35
Father's BMI (kg.m^{-2})	27,65	21,55	37,98	5,97
Mother's BMI (kg.m^{-2})	23,60	18,40	38,86	7,05
Child's BMI (kg.m^{-2})	18,95	13,94	32,24	6,07
Father's exercise (%)	71,4	-	-	-
Mother's exercise (%)	42,8	-	-	-
Father's PAL ($\text{Kcal.kg}^{-1}.\text{day}^{-1}$)	43,07	39,65	52,47	4,60
Mother's PAL ($\text{Kcal.kg}^{-1}.\text{day}^{-1}$)	42,67	38,43	50,18	3,83
Child's PAL ($\text{Kcal.kg}^{-1}.\text{day}^{-1}$)	42,90	40,51	45,08	1,79

The family income and the social status were not different between sedentary and active children. Social status mean was 5,0 and 5,14 for active and sedentary groups, respectively. According to the ANEP classification these families are included in the B1 and A2 status. This means that social status were not a intervenient variable in the children physical activity level.

Child's BMI and mother's BMI, in both groups, presented an ideal range. The reference values for BMI for children of this age is about $18,40 \text{ kg/m}^2$ (COLE, 2000) and for adults is $20-25 \text{ kg/m}^2$. However, the father's BMI was above this adult reference indicating an overweight status for both groups.

The physical activity data shows that there were not significant differences between the groups neither to the exercise practice nor to the PAL. In the sedentary group, mothers presented slighter superior values in the exercise adherence (75%) and PAL ($44,00 \text{ Kcal.kg}^{-1}.\text{day}^{-1}$) compared to the father's groups in the same sedentary group (62,5% and $40,87 \text{ Kcal.kg}^{-1}.\text{day}^{-1}$, respectively). On the other hand, in the active group, the better values of exercise practice and PAL were obtained by the fathers. Maybe fathers exert more influence in the children's lifestyles because in the active group the percentage of fathers that exercise (71,42%) was almost twice the percentage of mothers that regularly exercise (42,85%). To confirm this hypothesis it would be necessary a bigger sample.

Pearson's product moment correlation among all variables is presented in Table 3.

Table 3. Pearson's product moment correlation

	Family income	Social status	Father's BMI	Mother's BMI	Child's BMI	Father's exercise	Mother's exercise	Father's PAL	Mother's PAL	Child's PAL
Family income	1,00									
Social status	0,30	1,00								
Father's BMI	0,37	0,17	1,00							
Mother's BMI	-0,37	-0,01	-0,31	1,00						
Child's BMI	0,43	0,11	0,27	0,05	1,00					
Father's exercise	0,31	0,05	0,39	-0,15	-0,01	1,00				
Mother's exercise	0,17	0,00	-0,02	0,21	0,30	-0,29	1,00			
Father's PAL	-0,60*	-0,25	0,27	0,04	-0,26	-0,15	-0,31	1,00		
Mother's PAL	-0,34	-0,36	0,13	0,11	-0,24	0,32	-0,23	0,36	1,00	
Child's PAL	-0,48	-0,08	0,15	0,24	0,16	-0,19	0,35	0,25	-0,17	1,00

In the correlation results there was a negative result between father's PAL and family income, meaning that the bigger the income the smaller the fathers' PAL. Maybe families with high family income have a longer father's work time, or even because the facilities generated by technology decrease the necessity to perform physical tasks and consequently they decrease total energy expenditure as well.

There is a lack of previous studies to compare our findings because the majority of this kind of research only involves mothers or is made by using surveys to ask to children about the parents' exercise and incentive to children to exercise (WESTERSTAHL, et al., 2005, VILHJALMSSON, et al., 1998).

In study by Westerstahl, et al., (2005) there is a discussion about how far the complexity of variables determines the individual physical activity behavior and the authors point out the positive influence of parents and socioeconomic status in children PAL, such variables can influence children to take part in systematic sport since early childhood. Moreover, compared adolescents from different educational programs, Westerstahl and colleges, find that PAL differences can be explained by socioeconomic status, parents' support, perception of health and negative attitudes toward free physical activity (atividade física ao ar livre). Similar results were obtained by Vilhjalmsson, et al. (1998), in which parents' and familiar's emotional support and best friend physical activity showed strong relation to PAL in Irish adolescents.

In the present study there was no relationship between parents' PAL and child's PAL, which could mean that being active or sedentary is does not depend on the habitual parent's exercise participation. These data disagreed with the study by MOORE (1991) in which the child's PAL is influenced by parents' PAL.

The body composition result compared to PAL did not show significant results, meaning that the child's BMI is not related neither to the subject's PAL nor to the parents' BMI. Similar results were found by Treuth and colleges (2000) in which did not show relationship between energy expenditure or 8-to-9-years old girls and the parent's body composition (obese or not).

Other variables can be involved to the children lifestyle (sedentary or active). Steinbeck (2001) suggests that age, developmental status, gender, socioeconomic status, ethnic, parents being active or not, positive stimuli, aerobic fitness, obesity and genetics should be considered and the possibility of variance in such factors generates differences among children.

CONCLUSION

Socioeconomic status did not differ between the active and sedentary groups because the sample was selected from particular educational system. Moreover, the parents' scholarship, number of goods and family income was similar. Mother's BMI and child's BMI were according to the normal data, however father's BMI indicated overweight. The comparison between sedentary and active children, showed that BMI, exercise practice and parents' PAL did not differ significantly.

Except for the father's PAL and family income in this study, there was no relationship among BMI, socioeconomic status, exercise practice and habitual PAL between parents and their children, indicating existence of others intervenients variables.

One limitation of the present study is the small sample, which is an incentive for other researchers to study the factors that affect the children's lifestyle and what the contributions parents can bring to health and quality of life of their children.

REFERENCES

- BAR-OR, O. Juvenile obesity, physical activity and lifestyle changes. **The physician and sports medicine**. v. 28, n. 11, Nov. 2000.
- BERKEY, C. S.; ROCKETT, H.R.H.; FIELD, A. E.; GILLMAN, M.W.; FRAZIER, A. L.; CAMARGO, C. A.; COLDITZ, G. A. Activity, dietary intake and weight changes in a longitudinal study of preadolescent and adolescent boys and girls. **Pediatrics**. v. 105, n. 4, p.e56, 2000.
- BLAAK, E. E.; WESTERTERP, K. R.; BAR-OR, O.; WOUTERS, L. J. M.; SARIS, W. H. M. Total energy expenditure and spontaneous activity in relation to training in obese boys. **American journal of clinical nutrition**. v.55, p. 777-782, 1992.
- BOREHAM, C. A.; TWISK, J.; SAVAGE, M. J.; CRAN, G. W.; STRAIN, J. J. Physical activity, sports participation, and risk factors in adolescents. **Medicine and science in sports and exercise**. v. 29, n. 6, p. 788-793, 1997.
- BOUCHARD, C.; TREMBLAY, A.; LeBLANC, C.; LORTIE, G.; SAUARD, R.; THERIAULT, A. A Method to assess energy expenditure in children and adults. **American journal of clinical nutrition**. v. 37, p. 461-467, 1983.
- BUCHALLA, _____. Fontes do Projeto Estimativa dos Custos Atribuídos à Obesidade no Brasil, de Sichieri, R.; Vianna, C. M. e Coutinho, W. **Revista Veja**, 9Abr 2003.
- BRUM, V. P. C. A influência de diferentes níveis de atividade física e sexo sobre a aptidão aeróbica e anaeróbica de crianças pré-púberes da cidade de Curitiba-Pr. **Dissertação de Mestrado, Curso de Pós-graduação em Educação Física da Universidade Federal do Paraná**. 2004
- COLE, T. J. Establishing a standard definition for child overweight and obesity worldwide: international survey. **British medical journal**. v. 6, 2000.
- ELIAKIM, A.; SCHEETT, T.; ALLMENDINGER, N.; BRASEL, J. A.; COOPER, D. M. Training, muscle volume, and energy expenditure in nonobese American girls. **Journal of applied physiology**. v. 90, n. 1, p. 35-44, 2001.
- ENGESTRON, E. M.; ANJOS, L. A. Relação entre o estado nutricional materno e sobrepeso nas crianças brasileiras. **Revista Saúde Pública**. v. 30, n. 3, 1996.
- EPSTEIN, L. H.; PALUCH, R. A.; KALAKANIS, L. E.; GOLDFIELD, G. S.; CERNY, F. J.; ROEMMICH, J. N. How much activity do youth get? A quantitative review of heart rate measured activity. **Pediatrics**. v. 108, n. 3, p. e44, 2001.
- GUYTON, A. C.; HALL, J. E. **Tratado de Fisiologia Médica**. 9ª ed. RJ: Guanabara Koogan, 1997.
- IBGE Instituto Brasileiro de Geografia e Estatística, 2001. www.portal.saude.gov.br. Acessado em 2003.
- LAITINEN, J.; POWER, C. e JÄRVELIN, M. R. Family social class, maternal body mass index, childhood body mass index, and age at menarche as predictors of adult obesity. **American journal of clinical nutrition**. v. 74, n. 3, p. 287-294, 2001.
- MACHADO, H. S. A relação entre níveis de atividade física, aptidão física e os padrões motores fundamentais de escolares na faixa etária de 7 e 8 anos da cidade de Curitiba, Paraná. **Relatório técnico-científico apresentado ao programa institucional de bolsas de Iniciação Científica PIBIC/CNPq - Universidade Federal do Paraná**, 2002.
- MOORE, L. L. Influence of parents' physical activity levels on activity levels of young children. **J. Pediatrics**. v. 118, p. 215-219, 1991.
- OLIVEIRA, A. R. Fatores influenciadores na determinação do nível de aptidão física em crianças. **Revista Synopsis**. v. 7, p. 48-62, 1996.
- RIBEIRO, I. C. **Obesidade entre escolares da rede pública de ensino de Vila Mariana São Paulo: Estudo de caso-controle**. Dissertação de Mestrado USP/Escola Paulista de Medicina, 2001.
- ROSS, J. G. ; PATE, R.R. The national children and youth fitness study II: New health-related fitness norms. **Journal of physical education, recreation and dance**. v. 58, 1987.
- ROWLANDS, A. V.; ESTON, R. G.; INGLEDEW, D. K. Relationship between activity levels, aerobic fitness, and body fat in 8-to-10-year-old children. **Journal of applied physiology**. v. 86, n. 4, p. 1428-1435, 1999.
- SAELEN, B. E.; STEPHEN, R. D. **Childhood obesity: causes and therapies**. Current opinion in endocrinology e diabetes. v.10, n. 1, p. 3-8, 2003.
- STEINBECK, K. S. The importance of physical activity in the prevention of overweight and obesity in childhood: a review and an opinion. **Obesity reviews**. v. 2, p. 117-130, 2001.
- TREUTH, M. S; BUTTE, N. F.; WONG, W. W. Effects of familial predisposition to obesity on energy expenditure in multiethnic prepubertal girl. **American journal of clinical nutrition**. v. 71, p. 893-900, 2000.
- UKKOLA, O.; BOUCHARD, C. Fatores genéticos e obesidade infantil. **Anais Nestlé**. v. 62, p. 12-21, 2002.

VILHJALMSSON, R.; THORLINDSSON, T. Factors related to physical activity: a study of adolescents. **Soc. science medicine**. v. 47, n.5, p.665-675, 1998.
 VILLARES, S. M. F.; RIBEIRO, M. M.; SILVA, A. G. Obesidade infantil e exercício. **ABESO**. v. 13, p. 12-16, 2003.
 WESTERSTAHL, M.; BARNEKOW-BERGKVIST, M.; JANSSON, E. Low physical activity among adolescents in practical education. **Scandinavian journal medicine science**, 2005.

Contact Address:

Departamento de Educação Física - UFPR
 Centro de Pesquisa em Exercício e Esporte
 Endereço: Rua Coração de Maria, 92, Jd. Botânico, CEP 80-215-370
 Curitiba, Paraná Brasil
André Michelin 1@hotmail.com 9654-4512 (André)

RELATIONSHIP BETWEEN BMI, SOCIOECONOMIC STATUS, EXERCISE PARTICIPATION AND HABITUAL PHYSICAL ACTIVITY LEVEL OF FATHERS, MOTHERS AND THEIR CHILDREN
ABSTRACT

The purpose of this study is to establish the relationship between BMI, socioeconomic status, regular exercise practice and daily physical activity level of children and their parents from private schools of Curitiba-PR. The sample was composed of 45 subjects (15 eight-year-old school children, 15 fathers and 15 mothers). This sample was divided in two groups: sedentary children and their parents (n=24) and active children and their parents (n=21). At first, all subjects' body mass and stature were assessed, and then the BMI was calculated. A questionnaire was used to assess the socioeconomic status, and the regular exercise practice and habitual physical activity level was determined by using, from all subjects (fathers, mothers and children), a three-day diary (BOUCHARD *et al.*, 1983). Group comparisons were made by one-way ANOVA's and Pearson moment product was used to verify the relationship among all variables (P<0,05). The correlation results did not show significant difference between father's physical activity level and the family income (r=0,60), which means that the higher the family income the lower the father's PAL. In this sample, the fact of being either active or sedentary children is not related to the parents' physical activity. However, more studies, with large sample, are necessary to investigate this issue. **Key-words:** physical activity level, socioeconomic status, parents and children.

LA RELATION ENTRE LE IMC, LE NIVEAU ASSOCIÉ-ÉCONOMIQUE, LE EXERCICE PHYSIQUE ET LE NIVEAU D'ACTIVITÉ PHYSIQUE HABITUELLE DOS PÈRE, MÈRES ET LE FILS
RESUMÉE

Objectif: vérifier la relation entre le IMC, le niveau associé-économique, la pratiques des exercices physiques réguliers et le niveau d'activité physique habituelle et scolaire du filet particulier de l'éducation de Curitiba, Pr et ses pères respectifs. Methodologie: Le groupe été composé pour 45 individus, étant 15 scolaire avec 8 ans, 15 père et 15 mères. Le groupe se divise dans deux: sédentaire scolaire et ses pères (n=24) et actifs scolaire et ses pères (n=21). Ont été évalué masse corporelle, la stature et le IMC. Por la évaluation le niveau associé-économique et le pratique de l'exercice physique des pères ont été utilisées un entrevue et por la évaluation du niveau d'activité physique habituelle (NAFH) du père, la mère et le fils utilisées entrevue au sujet d'activité quotidienne (BOUCHARD *et al.*, 1983). Des analyses de variance (one-way) et corrélations de moment de Pearson ont été utilisées por l'analyse des résultats, avend un niveau alpha établie en 0,05. Résultant: la corrélation il était significative entre le NAFH du père et du revenu familial (r=-0,60), ce qui signifie que plus le revenu de famille est haut plus pal du père est inférieur. Dans cet échantillon, le fait de l'enfant actif ou sédentaire à être n'est pas relié le pratique d'activité physique des pères. Cependant, mais d'études sont nécessaires, depuis l'échantillon de cette étude a été limité dans le nombre d'individus. **Mots-clé:** niveau d'activité physique, niveau associé-économique, pères et fils

LA RELACIÓN ENTRE EL IMC, EL NIVEL SOCIO-ECONOMICO, EJERCICIO FISICO E EL NIVEL DE ACTIVIDAD FISICA HABITUAL DE LOS PADRES, MADRES E HIJOS.
RESUMEN

El objetivo de este estudio fue verificar la relación entre el IMC, el nivel sócio-económico, la practica habitual de ejercicio físico, el nivel de actividad física diaria de alumnos de la red particular de enseñanza en Curitiba, Paraná y el de sus respectivos padres. La muestra fue compuesta por 45 individuos, siendo 15 alumnos de 8 años de edad, 15 padres y 15 madres. Esta muestra fue dividida em 2 dos grupos: alumnos sedentários y sus padres (n=24) y alumnos activos y sus padres (n=21). Inicialmente se evaluaron la masa corporal y la estatura de todos los individuos obteniendo el IMC, para el nivel socio-económico e la práctica de ejercicio físico de los padres fue usado un cuestionario y para la evaluación del nivel de la actividad física habitual (NAFH) del padre, de la madre y del hijo fue usado um recordatorio propuesto por BOUCHARD *et al.*, (1983). Em la análisis estadística fueron calculados las medias y el desvio-patrón ANOVA (one-way) y Correlación de Pearson, com el nivel alpha de 0,05. En los resultados de la correlación hubo significación entre el NAFH del padre y el total de lo que gana la familia (r= - 0,60), o sea, cuanto mas grande el poder adquisitivo de la familia menos es el NAFH del padre. En esta muestra, el facto del nino ser o activo o sedentário no está relacionado a la práctica de actividad física de los padres. Entretanto, mas estudios son necesarios, visto que la muestra deste estudio fue restrita em número de indivíduos.

Palabras-Clave: nivel de actividad físico, nivel sócio-económico, padres y hijos

RELAÇÃO ENTRE O IMC, NÍVEL SÓCIO-ECONÔMICO, EXERCÍCIO FÍSICO E NÍVEL DE ATIVIDADE FÍSICA HABITUAL DOS PAIS, MÃES E FILHOS
RESUMO

Objetivo: verificar a relação entre o IMC, o nível sócio-econômico, a prática de exercícios físicos regulares e o nível de atividade física diário dos escolares da rede particular de ensino de Curitiba/Pr e de seus respectivos pais. Métodos: amostra foi composta por 45 indivíduos, sendo 15 escolares com 8 anos, 15 pais e 15 mães, divididas em dois grupos: escolares sedentários e seus pais (n=24) e escolares ativos e seus pais (n=21). Inicialmente avaliou-se a massa corporal e a estatura obtendo-se o IMC, para o nível sócio-econômico e a prática de exercício físico dos pais foi utilizado um questionário e para avaliação do nível de atividade física habitual (NAFH) do pai, da mãe e do filho (a) utilizou-se um recordatório (BOUCHARD *et al.*, 1983). Na análise estatística foram calculados as médias e desvio-padrão, ANOVA (one-way) e Correlação de Pearson, com nível *alpha* de 0,05. Resultados: na correlação houve significância entre o NAFH do pai e a renda familiar (r=-0,60), ou seja, quanto maior o poder aquisitivo da família menor é o NAFH do pai. Conclusão: o fato da criança ser ativa ou sedentária não está relacionado a prática de atividade física dos pais. Entretanto, mais estudos são necessários, visto que a amostra deste estudo foi restrita no número de indivíduos.

Palavras-chave: nível de atividade física, nível sócio-econômico, pais e filhos.